

HELP PROTECT YOUR DOG YEAR-ROUND.

HEARTGARD Plus offers trusted heartworm disease prevention, plus broad-spectrum treatment and control against hookworms and roundworms – ALL in the exclusive Real-Beef Chewable that dogs love to take⁶ and owners prefer to give.⁷



IMPORTANT SAFETY INFORMATION: HEARTGARD Plus is well tolerated. All dogs should be tested for heartworm infection before starting a preventive program. Following the use of HEARTGARD Plus, digestive and neurological side effects have rarely been reported. For more information, contact your veterinarian or visit www.HEARTGARD.com.

EASY TO GIVE. PROTECTION TO LIVE.

TRUSTED, EFFECTIVE, MONTHLY PREVENTION


It's up to you to help prevent heartworm disease and control hookworms and roundworms in your pet – and it's as easy as the Real-Beef Chewable your dog will love to take!⁶

THE #1 CHOICE
OF VETERINARIANS.⁸



Heartgard[®]
(ivermectin/pyrantel) **Plus**

Go to HEARTGARD.com for details
and to sign up for FREE monthly reminders.



Heartworm disease
prevention is easy with

Heartgard[®]
(ivermectin/pyrantel) **Plus**



HEARTWORMS— A REAL & PRESENT DANGER

HOW DOES IT WORK?

When you give HEARTGARD® Plus on the 1st of the month, it takes care of any heartworm infection picked up in the last 30 days.⁵ HEARTGARD Plus should be given EVERY 30 DAYS to help prevent heartworm disease in your dog.

JUNE

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

IMPORTANT SAFETY INFORMATION:

HEARTGARD Plus is well tolerated. All dogs should be tested for heartworm infection before starting a preventive program. Following the use of HEARTGARD Plus, digestive and neurological side effects have rarely been reported. For more information, contact your veterinarian or visit www.HEARTGARD.com.

1
MILLION

Dogs infected with adult heartworms in the U.S.¹



Mosquitoes transmit heartworm larvae from infected dogs to healthy dogs.



Heartworms live in the hearts and lungs of infected dogs.



Common signs of infection include coughing, difficulty breathing and sluggishness. Early infections may show no signs.



Untreated heartworm infections can be fatal.



Other carriers include wolves, foxes, ferrets and coyotes.

INTESTINAL PARASITES 101


All dogs are at risk for hookworms and roundworms – dangerous parasites that can spread to people, too.

HOOKWORMS – A BURDEN FOR PETS AND PEOPLE



IN DOGS:


- Live in dogs' intestines
- Can cause anemia, debilitation, and even death.



IN HUMANS:


- Enter through the skin or accidental ingestion
- Can cause rashes, itching, eruptions through the skin and even organ inflammation.²

ROUNDWORMS – BEYOND THE “YUCK” FACTOR FOR THE WHOLE FAMILY



IN DOGS:

- Can infect over 90% of puppies under 3 months old.³
- Can cause diarrhea, weight loss and swollen abdomen



IN HUMANS:

- The CDC reports 10,000 people in the U.S. per year with roundworm infections.⁴
- Can cause impaired – or loss of – vision.⁴

ONE MOSQUITO IS ALL IT TAKES TO TRANSMIT HEARTWORMS TO YOUR DOG.

All dogs are at risk for heartworm disease no matter where they live!
Once a dog is infected with heartworms, treatment can be expensive, difficult and unsuccessful.

Heartgard[®] Plus
(ivermectin/pyrantel)

CHEWABLES

CAUTION: Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.

INDICATIONS: For use in dogs to prevent canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for a month (30 days) after infection and for the treatment and control of ascarids (*Toxocara canis*, *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*, *Ancylostoma braziliense*).

DOSAGE: HEARTGARD[®] Plus (ivermectin/pyrantel) should be administered orally at monthly intervals at the recommended minimum dose level of 6 mcg of ivermectin per kilogram (2.72 mcg/lb) and 5 mg of pyrantel (as pamoate salt) per kg (2.27 mg/lb) of body weight. The recommended dosing schedule for prevention of canine heartworm disease and for the treatment and control of ascarids and hookworms is as follows:

Dog Weight	Chewables Per Month	Ivermectin Content	Pyrantel Content	Color Coding On Foil Backing and Carton
Up to 25 lb	1	68 mcg	57 mg	Blue
26 to 50 lb	1	136 mcg	114 mg	Green
51 to 100 lb	1	272 mcg	227 mg	Brown

HEARTGARD Plus is recommended for dogs 6 weeks of age and older.
For dogs over 100 lb use the appropriate combination of these chewables.

ADMINISTRATION: Remove only one chewable at a time from the foil-backed blister card. Return the card with the remaining chewables to its box to protect the product from light. Because most dogs find HEARTGARD Plus palatable, the product can be offered to the dog by hand. Alternatively, it may be added intact to a small amount of dog food. The chewable should be administered in a manner that encourages the dog to chew, rather than to swallow without chewing. Chewables may be broken into pieces and fed to dogs that normally swallow treats whole.

Care should be taken that the dog consumes the complete dose, and treated animals should be observed for a few minutes after administration to ensure that part of the dose is not lost or rejected. If it is suspected that any of the dose has been lost, redosing is recommended.

HEARTGARD Plus should be given at monthly intervals during the period of the year when mosquitoes (vectors), potentially carrying infective heartworm larvae, are active. The initial dose must be given within a month (30 days) after the dog's first exposure to mosquitoes. The final dose must be given within a month (30 days) after the dog's last exposure to mosquitoes.

When replacing another heartworm preventive product in a heartworm disease preventive program, the first dose of HEARTGARD Plus must be given within a month (30 days) of the last dose of the former medication.

If the interval between doses exceeds a month (30 days), the efficacy of ivermectin can be reduced. Therefore, for optimal performance, the chewable must be given once a month on or about the same day of the month. If treatment is delayed, whether by a few days or many, immediate treatment with HEARTGARD Plus and resumption of the recommended dosing regimen will minimize the opportunity for the development of adult heartworms.

Monthly treatment with HEARTGARD Plus also provides effective treatment and control of ascarids (*T. canis*, *T. leonina*) and hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*). Clients should be advised of measures to be taken to prevent reinfection with intestinal parasites.

EFFICACY: HEARTGARD Plus Chewables, given orally using the recommended dose and regimen, are effective against the tissue larval stage of *D. immitis* for a month (30 days) after infection and, as a result, prevent the development of the adult stage. HEARTGARD Plus Chewables are also effective against canine ascarids (*T. canis*, *T. leonina*) and hookworms (*A. caninum*, *U. stenocephala*, *A. braziliense*).

ACCEPTABILITY: In acceptability and field trials, HEARTGARD Plus was shown to be an acceptable oral dosage form that was consumed at first offering by the majority of dogs.

PRECAUTIONS: All dogs should be tested for existing heartworm infection before starting treatment with HEARTGARD Plus which is not effective against adult *D. immitis*. Infected dogs must be treated to remove adult heartworms and microfilariae before initiating a program with HEARTGARD Plus.

While some microfilariae may be killed by the ivermectin in HEARTGARD Plus at the recommended dose level, HEARTGARD Plus is not effective for microfilariae clearance. A mild hypersensitivity-type reaction, presumably due to dead or dying microfilariae and particularly involving a transient diarrhea, has been observed in clinical trials with ivermectin alone after treatment of some dogs that have circulating microfilariae.

Keep this and all drugs out of the reach of children.

In case of ingestion by humans, clients should be advised to contact a physician immediately. Physicians may contact a Poison Control Center for advice concerning cases of ingestion by humans.

Store between 68°F - 77°F (20°C - 25°C). Excursions between 59°F - 86°F (15°C - 30°C) are permitted. Protect product from light.

ADVERSE REACTIONS: In clinical field trials with HEARTGARD Plus, vomiting or diarrhea within 24 hours of dosing was rarely observed (1.1% of administered doses). The following adverse reactions have been reported following the use of HEARTGARD: Depression/lethargy, vomiting, anorexia, diarrhea, mydriasis, ataxia, staggering, convulsions and hypersalivation.

SAFETY: HEARTGARD Plus has been shown to be bioequivalent to HEARTGARD, with respect to the bioavailability of ivermectin. The dose regimens of HEARTGARD Plus and HEARTGARD are the same with regard to ivermectin (6 mcg/kg). Studies with ivermectin indicate that certain dogs of the Collie breed are more sensitive to the effects of ivermectin administered at elevated dose levels (more than 16 times the target use level) than dogs of other breeds. At elevated doses, sensitive dogs showed adverse reactions which included mydriasis, depression, ataxia, tremors, drooling, paresis, recumbency, excitability, stupor, coma and death. HEARTGARD demonstrated no signs of toxicity at 10 times the recommended dose (60 mcg/kg) in sensitive Collies. Results of these trials and bioequivalency studies, support the safety of HEARTGARD products in dogs, including Collies, when used as recommended.

HEARTGARD Plus has shown a wide margin of safety at the recommended dose level in dogs, including pregnant or breeding bitches, stud dogs and puppies aged 6 or more weeks. In clinical trials, many commonly used flea collars, dips, shampoos, anthelmintics, antibiotics, vaccines and steroid preparations have been administered with HEARTGARD Plus in a heartworm disease prevention program.

In one trial, where some pups had parvovirus, there was a marginal reduction in efficacy against intestinal nematodes, possibly due to a change in intestinal transit time.

HOW SUPPLIED: HEARTGARD Plus is available in three dosage strengths (See DOSAGE section) for dogs of different weights. Each strength comes in convenient cartons of 6 and 12 chewables.

For customer service, please contact Merial at 1-888-637-4251.

IMPORTANT SAFETY INFORMATION:

HEARTGARD Plus is well tolerated. All dogs should be tested for heartworm infection before starting a preventive program. Following the use of HEARTGARD Plus, digestive and neurological side effects have rarely been reported. For more information, contact your veterinarian or visit www.HEARTGARD.com.

- Weyenberg K. Team up against heartworm. *The Team*. 2009. 6(4)10-12.
- Centers for Disease Control and Prevention. What Every Pet Owner Should Know About Roundworms & Hookworms. Available at <http://stacks.cdc.gov/view/cdc/13560>. Accessed December 16, 2014.
- Schantz PM. Zoonotic ascarids and hookworms: the role for veterinarians in preventing human disease. In: *Emerging Vector-Borne and Zoonotic Diseases, The Compendium Suppl* 2002;24(1):47-52.
- Roundworm Infection, Toxocariasis, FAST FACTS. Available at http://www.cfsph.iastate.edu/FastFacts/pdfs/toxocariasis_F.pdf. Accessed December 16, 2014.
- Campbell WC. Use of ivermectin in dogs and cats. In: Campbell WC, ed. *Ivermectin and Abamectin*. New York: Springer-Verlag, 1989:246-248.
- Freedom of Information: NADA140-971 (January 15, 1993).
- Vet-dispensed; Heartworm Preventives Pet Owner Awareness and Attitude Tracking August 2014. Bruno and Ridgway. Data on file at Merial.
- Vet-dispensed; CMI data, September 2014; data on file at Merial.



Merial is now part of Boehringer Ingelheim.

©HEARTGARD and the Dog & Hand logo are registered trademarks of Merial. ©2017 Merial, Inc., Duluth, GA. All rights reserved. HGD16BROCHURE (12/16).

